

Actins

Franke W.W., Stehr S., Stumpp S., Kuhn C., Heid H., Rackwitz H.-R., Schnölzer M., Baumann R., Holzhausen H.-J., Moll R.

Specific immunohistochemical detection of cardiac/fetal α -actin in human cardiomyocytes and regenerating skeletal muscle cells, 60:245–250

Hayward S.W., Brody J.R., Cunha G.R.

An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219–227

Lincz L.F., Buret A., Burns G.F.

Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216–274

Adipose differentiation

Casimir D.A., Wilson-Miller C., Ntambi J.M.

Preadipocyte differentiation blocked by prostaglandin stimulation of prostanoid FP₂ receptor in murine 3T3-L1 cells, 60:203–210

Ntambi J.M., Takova T.

Role of Ca²⁺ in the early stages of murine adipocyte differentiation as evidenced by calcium mobilizing agents, 60:151–158

Acute lymphoblastic leukemia

Maki A., Mohammad R.M., Smith M., Al-Katib A.

Role of ubiquitin carboxyl terminal hydrolase in the differentiation of human acute lymphoblastic leukemia cell line, Reh, 60:59–66

Amphibia

Izutsu Y., Yoshizato K., Tochinali S.

Adult-type splenocytes of *Xenopus* induce apoptosis of histocompatible larval tail cells in vitro, 60:277–286

Lopashov G.V., Golubeva O.N., Zviadadze K.G.

Inductive capacity of living eye tissues from adult frogs, 61:237–242

Okamoto M.

Simultaneous demonstration of lens regeneration from dorsal iris and tumour production from ventral iris in the same newt eye after carcinogen administration, 61:285–292

Zilakos N.P., Zafiratos C.S., Parchment R.E.

Stage-dependent genetically-based deformities of the regenerating newt limb from 4-nitroquinoline-N-oxide mutagenesis: Potential embryonic regulation of cancer, 60:67–74

Androgenic ES cells

McKarney L.A., Overall M.L., Dziadek M.

Expression of H19 and Igf2 genes in uniparental mouse ES cells during in vitro and in vivo differentiation, 60:75–86

Antisense oligodeoxynucleotides

Maki A., Mohammad R.M., Smith M., Al-Katib A.

Role of ubiquitin carboxyl terminal hydrolase in the differentiation of human acute lymphoblastic leukemia cell line, Reh, 60:59–66

Aorta

Slomp J., Gittenberger-de Groot A.C., Kotliansky V.E.,

Glukhova M.A., Bogers A.J.J.C., Poelman R.E.

Cytokeratin expression in human arteries pertinent to intimal thickening formation in the ductus arteriosus, 61:305–311

Apoptosis

Izutsu Y., Yoshizato K., Tochinali S.

Adult-type splenocytes of *Xenopus* induce apoptosis of histocompatible larval tail cells in vitro, 60:277–286

Maki A., Mohammad R.M., Smith M., Al-Katib A.

Role of ubiquitin carboxyl terminal hydrolase in the differentiation of human acute lymphoblastic leukemia cell line, Reh, 60:59–66

Arachidonic acid

Casimir D.A., Wilson-Miller C., Ntambi J.M.

Preadipocyte differentiation blocked by prostaglandin stimulation of prostanoid FP₂ receptor in murine 3T3-L1 cells, 60:203–210

Armadillo proteins

Paffenholz R., Franke W.W.

Identification and localization of a neuronally expressed member of the plakoglobin/armadillo multigene family, 61:293–304

Aromatase

Dorizzi M., Richard-Mercier N., Pieau C.

The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193–201

Atrial natriuretic peptide

Koide M., Akins R.E., Harayama H., Yasui K., Yokota M., Tuan R.S.

Atrial natriuretic peptide accelerates proliferation of chick embryonic cardiomyocytes in vitro, 61:1–11

Auxin

Meyer A.D., Aebi R., Meins Jr. F.

Tobacco plants carrying a tms locus of Ti-plasmid origin and the *HI-1* allele are tumor prone, 61:213–221

Basement membrane

Breitkreutz D., Stark H.-J., Mirancea N., Tomakidi P.,

Steinbauer H., Fusenig N.E.

Integrin and basement membrane normalization in mouse grafts of human keratinocytes - Implications for epidermal homeostasis, 61:195–209

Basic fibroblast growth factor

Toriyama K., Muramatsu H., Hoshino T., Torii S., Muramatsu T.

Evaluation of heparin-binding growth factors in rescuing morphogenesis of heparitinase-treated mouse embryonic lung explants, 61:161–167

B-cell

Maki A., Mohammad R.M., Smith M., Al-Katib A.

Role of ubiquitin carboxyl terminal hydrolase in the differentiation of human acute lymphoblastic leukemia cell line, Reh, 60:59–66

Beta-cell of pancreas

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F.,

Jami J., Buccini D.

Human insulin gene expression in transgenic mice: mutational analysis of regulatory region, 60:309–316

Blastocoelomic antigens

Vafa O., Goetzl L., Poccia D., Nishioka D.
Localization and characterization of blastocoelomic extracellular matrix antigens in early sea urchin embryos and evidence for their proteolytic modification during gastrulation, 60:129-138

Breast cancer cells

Jing Y., Zhang J., Waxman S., Mira-y-Lopez R.
Upregulation of cytokeratins 8 and 18 in human breast cancer T47D cells is retinoid-specific and retinoic acid receptor-dependent, 60:109-117

Brefeldin A

Chazaud B., Muriel M.-P., Aubery M., Decastel M.
Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Cadherins

Anzai H., Kitadai Y., Bucana C.D., Sanchez R., Omoto R., Fidler I.J.
Intratumoral heterogeneity and inverse correlation between expression of E-cadherin and collagenase type IV in human gastric carcinomas, 60:119-127

Eng H., Herrenknecht K., Semb H., Starzinski-Powitz A., Ringertz N., Gullberg D.
Effects of divalent cations on M-cadherins expression and distribution during primary rat myogenesis in vitro, 61:169-176
Liebert M., Hubbel A., Chung M., Wedemeyer G., Lomax M.I., Hegeman A., Yuan T.Y., Brozovich M., Wheelock M.J., Grossman H.B.
Expression of *mal* is associated with urothelial differentiation in vitro: identification by differential display reverse-transcriptase polymerase chain reaction, 61:177-185

Schäfer S., Stumpp S., Franke W.W.
Immunological identification and characterization of the desmosomal cadherin Dsg2 in coupled and uncoupled epithelial cells and in human tissues, 60:99-108

Calcineurin

Horn F., Gross J.
A role for calcineurin in *Dictyostelium discoideum* development, 60:269-275

Calcium

Horn F., Gross J.
A role for calcineurin in *Dictyostelium discoideum* development, 60:269-275

Lincz L.F., Buret A., Burns G.F.
Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216-274

Ntambi J.M., Takova T.
Role of Ca^{2+} in the early stages of murine adipocyte differentiation as evidenced by calcium mobilizing agents, 60:151-158

Eng H., Herrenknecht K., Semb H., Starzinski-Powitz A., Ringertz N., Gullberg D.
Effects of divalent cations on M-cadherins expression and distribution during primary rat myogenesis in vitro, 61:169-176

cAMP induction

Yang C., Brar S.K., Desbarats L., Siu C.-H.
Synthesis of the Ca^{2+} -dependent cell adhesion molecule DdCAD-1 is regulated by multiple factors during *Dictyostelium* development, 61:275-284

Cancer therapy

Naka K., Yokozaki H., Domen T., Hayashi K., Kuniyasu H., Yasui W., Lotan R., Tahara E.
Growth inhibition of cultured human gastric cancer cells by 9-*cis*-retinoic acid with induction of cdk inhibitor Waf1/Cip1/Sdi1/p21 protein, 61:313-320

Carassius

Marí-Beffa M., Fernandez-Llebrez J.A.S.P., Becerra J.
Histochemically defined cell states during tail fin regeneration in teleost fishes, 60:139-149

Carbamoylphosphate synthetase I

Koike T., Shiojiri N.
Differentiation of the mouse hepatic primordium cultured in vitro, 61:35-43

Cardiomyocytes

Franke W.W., Stehr S., Stumpp S., Kuhn C., Heid H., Rackwitz H.-R., Schnölzer M., Baumann R., Holzhausen H.-J., Moll R.
Specific immunohistochemical detection of cardiac/fetal α -actin in human cardiomyocytes and regenerating skeletal muscle cells, 60:245-250

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F., Jami J., Buccini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316

Koide M., Akins R.E., Harayama H., Yasui K., Yokota M., Tuan R.S.
Atrial natriuretic peptide accelerates proliferation of chick embryonic cardiomyocytes in vitro, 61:1-11

CAT (chloramphenicol acetyltransferase) expression

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F., Jami J., Buccini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316

Catalase

Radaeva S., Bannasch P.
Changes in catalase and glucose-6-phosphatase distribution patterns within oval cell compartment as possible differentiation markers during viral hepatocarcinogenesis in woodchucks, 60:169-178

Catenins

Paffenholz R., Franke W.W.
Identification and localization of a neuronally expressed member of the plakoglobin/armadillo multigene family, 61:293-304

CD44 isoforms

Weber B., Rösel M., Arch R., Möller P., Zöller M.
Transient expression of CD44 variant isoforms in the ontogeny of the rat: Ectoderm-, endoderm- and mesoderm-derived cells express different exon combinations, 60:17-29

Cell adhesion

Kadmon G., Altevogt P.
The cell adhesion molecule L1: species and cell-type-dependent multiple binding mechanisms, 61:143-150

Yang C., Brar S.K., Desbarats L., Siu C.-H.
Synthesis of the Ca^{2+} -dependent cell adhesion molecule DdCAD-1 is regulated by multiple factors during *Dictyostelium* development, 61:275-284

Cell communication

Matic M., Petrov I.N., Chen S., Wang C., Dimitrijevic S.D., Wolosin J.M.
Stem cells of the corneal epithelium lack connexins and metabolite transfer capacity, 61:251-260

Cell rearrangement

Lincz L.F., Buret A., Burns G.F.
Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216-274

CFU-S

Lin Y.I.

Day 11 mouse fetal thymus: phenotype and search for the point of commitment, 61:53-65

Chemoreceptors

Zhang C., Oakley B.

The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Chick embryos

Koide M., Akins R.E., Harayama H., Yasui K., Yokota M., Tuan R.S.

Atrial natriuretic peptide accelerates proliferation of chick embryonic cardiomyocytes in vitro, 61:1-11

Chondroblasts

Cohen-Tanugi A., Bolle M., Boy-Lefèvre M.-L., Anagnostou F., Forest N.

Phenotypic characterization of the 3/A/1D-1M osteogenic cell line derived from in vivo transplantation of 3/A/1D-1 chondroprogenitor murine teratocarcinoma cells, 60:327-337

Chondrocytes

Grigoriadis A.E., Heersche J.N.M., Aubin J.E.

Analysis of chondroprogenitor frequency and cartilage differentiation in a novel family of clonal chondrogenic rat cell lines, 60:299-307

Chondrogenesis

Cohen-Tanugi A., Bolle M., Boy-Lefèvre M.-L., Anagnostou F., Forest N.

Phenotypic characterization of the 3/A/1D-1M osteogenic cell line derived from in vivo transplantation of 3/A/1D-1 chondroprogenitor murine teratocarcinoma cells, 60:327-337

Grigoriadis A.E., Heersche J.N.M., Aubin J.E.

Analysis of chondroprogenitor frequency and cartilage differentiation in a novel family of clonal chondrogenic rat cell lines, 60:299-307

Wroblewski J., Engström M., Edwall-Arvidsson C., Sjöberg G., Sejersen T., Lendahl U.

Distribution of nestin in the developing mouse limb bud in vivo and in micro-mass cultures of cells isolated from limb buds, 61:151-159

Clonal cultures

Baroffio A., Hamann M., Bernheim L., Bochaton-Piallat M.-L., Gabbiani G., Bader C.R.

Identification of self-renewing myoblasts in the progeny of single human muscle satellite cells, 60:47-57

Collagen

Yuan S., Rosenberg L., Paraskevas S., Agapitos D., Duguid W.P. Transdifferentiation of human islets to pancreatic ductal cells in collagen matrix culture, 61:67-75

Collagenase IV

Anzai H., Kitadai Y., Bucana C.D., Sanchez R., Omoto R., Fidler I.J.

Intratumoral heterogeneity and inverse correlation between expression of E-cadherin and collagenase type IV in human gastric carcinomas, 60:119-127

Colon carcinoma

Lincz L.F., Buret A., Burns G.F.

Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216-274

Colorectal epithelium

Henderson K., Kirkland S.C.

Multilineage differentiation of cloned HRA-19 cells in serum-free medium: A model of human colorectal epithelial differentiation, 60:259-268

Conjunctival epithelium

Moyer P.D., Kaufman A.H., Zhang Z., Kao C.W.-C., Spaulding A.G., Kao W.W.-Y.

Conjunctival epithelial cells can resurface denuded cornea, but do not transdifferentiate to express cornea-specific keratin 12 following removal of limbal epithelium in mouse, 60:31-38

Connexins

Matic M., Petrov I.N., Chen S., Wang C., Dimitrijevic S.D., Wolosin J.M.

Stem cells of the corneal epithelium lack connexins and metabolite transfer capacity, 61:251-260

Corneal epithelium

Matic M., Petrov I.N., Chen S., Wang C., Dimitrijevic S.D., Wolosin J.M.

Stem cells of the corneal epithelium lack connexins and metabolite transfer capacity, 61:251-260

Moyer P.D., Kaufman A.H., Zhang Z., Kao C.W.-C., Spaulding A.G., Kao W.W.-Y.

Conjunctival epithelial cells can resurface denuded cornea, but do not transdifferentiate to express cornea-specific keratin 12 following removal of limbal epithelium in mouse, 60:31-38

Crown gall tumor

Meyer A.D., Aebi R., Meins Jr. F.

Tobacco plants carrying a *tms* locus of Ti-plasmid origin and the *HI-1* allele are tumor prone, 61:213-221**Cytokeratins**

Chazaud B., Muriel M.-P., Aubery M., Decastel M.

Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Hayward S.W., Brody J.R., Cunha G.R.

An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219-227

Jing Y., Zhang J., Waxman S., Mira-y-Lopez R.

Upregulation of cytokeratins 8 and 18 in human breast cancer T47D cells is retinoid-specific and retinoic acid receptor-dependent, 60:109-117

Plancha C.E.

Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Slomp J., Gittenberger-de Groot A.C., Kotliansky V.E., Glukhova M.A., Bogers A.J.J.C., Poelman R.E.

Cytokeratin expression in human arteries pertinent to intimal thickening formation in the ductus arteriosus, 61:305-311

Zhang C., Oakley B.

The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Cytoskeleton

Chazaud B., Muriel M.-P., Aubery M., Decastel M.

Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Plancha C.E.

Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Wride M.A.

Cellular and molecular features of lens differentiation: a review of recent advances, 61:77-93

Cytostatic effect

Naka K., Yokozaki H., Domen T., Hayashi K., Kuniyasu H., Yasui W., Lotan R., Tahara E.

Growth inhibition of cultured human gastric cancer cells by 9-*cis*-retinoic acid with induction of cdk inhibitor Waf1/Cip1/Sd1/p21 protein, 61:313-320

Dermal bone

Marí-Beffa M., Fernandez-Lliebrez J.A.S.P., Becerra J.

Histochemically defined cell states during tail fin regeneration in teleost fishes, 60:139-149

Dermal papilla

Matzuzaki T., Inamatsu M., Yoshizato K.

The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297

Dermal sheath

Matzuzaki T., Inamatsu M., Yoshizato K.

The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297

Desmin

Wroblewski J., Engström M., Edwall-Arvidsson C., Sjöberg G., Sejersen T., Lendahl U.

Distribution of nestin in the developing mouse limb bud in vivo and in micro-mass cultures of cells isolated from limb buds, 61:151-159

Desmogleins

Schäfer S., Stumpp S., Franke W.W.

Immunological identification and characterization of the desmosomal cadherin Dsg2 in coupled and uncoupled epithelial cells and in human tissues, 60:99-108

Desmosomes

Liebert M., Hubbel A., Chung M., Wedemeyer G., Lomax M.I., Hegeman A., Yuan T.Y., Brozovich M., Wheelock M.J., Grossman H.B.

Expression of *mal* is associated with urothelial differentiation in vitro: identification by differential display reverse-transcriptase polymerase chain reaction, 61:177-185

Schäfer S., Stumpp S., Franke W.W.

Immunological identification and characterization of the desmosomal cadherin Dsg2 in coupled and uncoupled epithelial cells and in human tissues, 60:99-108

Development

Durham P.L., Snyder J.M.

Regulation of the β 2 subunit chain of laminin in developing rabbit fetal lung tissue, 60:229-243

Horn F., Gross J.

A role for calcineurin in *Dictyostelium discoideum* development, 60:269-275

Paranko J., Seitz J., Meinhardt A.

Developmental expression of heat shock protein 60 (HSP60) in the rat testis and ovary, 60:159-167

Wride M.A.

Cellular and molecular features of lens differentiation: a review of recent advances, 61:77-93

Dexamethasone

Koike T., Shiojiri N.

Differentiation of the mouse hepatic primordium cultured in vitro, 61:35-43

Dictyostelium

Horn F., Gross J.

A role for calcineurin in *Dictyostelium discoideum* development, 60:269-275

Robinson V., Williams J.

A marker of terminal stalk cell terminal differentiation in *Dictyostelium*, 61:223-228

Schaap P., Tang Y., Othmer H.G.

A model for pattern formation in *Dictyostelium discoideum*, 60:1-16, 61:141

Yang C., Brar S.K., Desbarats L., Siu C.-H.

Synthesis of the Ca^{2+} -dependent cell adhesion molecule DdCAD-1 is regulated by multiple factors during *Dictyostelium* development, 61:275-284

Differentiation inducing factor (DIF)

Schaap P., Tang Y., Othmer H.G.

A model for pattern formation in *Dictyostelium discoideum*, 60:1-16, 61:141

Drosophila

Hess, K.A., Simone A.A., Petschek J.P.

Spatial and temporal expression of *4f-rnp* gene in *Drosophila melanogaster*, 61:103-111

Ductus arteriosus

Slomp J., Gittenberger-de Groot A.C., Koteliensky V.E.,

Glukhova M.A., Bogers A.J.J.C., Poelman R.E.

Cytokeratin expression in human arteries pertinent to intimal thickening formation in the ductus arteriosus, 61:305-311

Ectoderm

Zhang C., Oakley B.

The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Lopashov G.V., Golubeva O.N., Zviadadze K.G.

Inductive capacity of living eye tissues from adult frogs, 61:237-242

Electron microscopy

Radaeva S., Bannasch P.

Changes in catalase and glucose-6-phosphatase distribution patterns within oval cell compartment as possible differentiation markers during viral hepatocarcinogenesis in woodchucks, 60:169-178

Embryonic regulation

Zilakos N.P., Zafiratos C.S., Parchment R.E.

Stage-dependent genetically-based deformities of the regenerating newt limb from 4-nitroquinoline-N-oxide mutagenesis: Potential embryonic regulation of cancer, 60:67-74

Endochondral bone formation

Cohen-Tanugi A., Bolle M., Boy-Lefèvre M.-L., Anagnostou F., Forest N.

Phenotypic characterization of the 3A/1D-1M osteogenic cell line derived from in vivo transplantation of 3A/1D-1 chondroprogenitor murine teratocarcinoma cells, 60:327-337

Endoderm

Zhang C., Oakley B.

The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Endoplasmic reticulum

Chazaud B., Muriel M.-P., Aubery M., Decastel M.
Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Enterocytes

Chazaud B., Muriel M.-P., Aubery M., Decastel M.
Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Henderson K., Kirkland S.C.
Multilineage differentiation of cloned HRA-19 cells in serum-free medium: A model of human colorectal epithelial differentiation, 60:259-268

Lincz L.F., Buret A., Burns G.F.
Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216-274

Epidermis

Bickenbach J.R., Longley M.A., Bundman D.S., Dominey A.M., Bowden P.E., Rothnagel A.M., Roop D.R.
A transgenic mouse model that recapitulates the clinical features of both neonatal and adult forms of the skin disease epidermolytic hyperkeratosis, 61:129-139

Breitbart D., Stark H.-J., Mirancea N., Tomakidi P., Steinbauer H., Fusenig N.E.
Integrin and basement membrane normalization in mouse grafts of human keratinocytes - Implications for epidermal homeostasis, 61:195-209

Matzuzaki T., Inamatus M., Yoshizato K.
The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297

Epidermolytic hyperkeratosis (EHK)

Bickenbach J.R., Longley M.A., Bundman D.S., Dominey A.M., Bowden P.E., Rothnagel A.M., Roop D.R.
A transgenic mouse model that recapitulates the clinical features of both neonatal and adult forms of the skin disease epidermolytic hyperkeratosis, 61:129-139

Epithelia

Cella N., Cornejo-Urbe R.R., Montes G.S., Hynes N.E.,
The lysosomal-associated membrane protein LAMP-1 is a novel differentiation marker for HC11 mouse mammary epithelial cells, 61:113-120

Hayward S.W., Brody J.R., Cunha G.R.
An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219-227

Henderson K., Kirkland S.C.
Multilineage differentiation of cloned HRA-19 cells in serum-free medium: A model of human colorectal epithelial differentiation, 60:259-268

Lincz L.F., Buret A., Burns G.F.
Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216-274

Moyer P.D., Kaufman A.H., Zhang Z., Kao C.W.-C., Spaulding A.G., Kao W.W.-Y.
Conjunctival epithelial cells can resurface denuded cornea, but do not transdifferentiate to express cornea-specific keratin 12 following removal of limbal epithelium in mouse, 60:31-38

Schäfer S., Stumpp S., Franke W.W.
Immunological identification and characterization of the desmosomal cadherin Dsg2 in coupled and uncoupled epithelial cells and in human tissues, 60:99-108

Extracellular matrix

Durham P.L., Snyder J.M.
Regulation of the β 2 subunit chain of laminin in developing rabbit fetal lung tissue, 60:229-243

Mari-Beffa M., Fernandez-Llebrez J.A.S.P., Becerra J.
Histochemically defined cell states during tail fin regeneration in teleost fishes, 60:139-149

Vafa O., Goetzl L., Poccia D., Nishioka D.
Localization and characterization of blastocoelic extracellular matrix antigens in early sea urchin embryos and evidence for their proteolytic modification during gastrulation, 60:129-138

Wride M.A.
Cellular and molecular features of lens differentiation: a review of recent advances, 61:77-93

Eye

Lopashov G.V., Golubeva O.N., Zviadadze K.G.
Inductive capacity of living eye tissues from adult frogs, 61:237-242

Fetal lung

Durham P.L., Snyder J.M.
Regulation of the β 2 subunit chain of laminin in developing rabbit fetal lung tissue, 60:229-243

Fiber type

Rosenblatt J.D., Parry D.J., Partridge T.A.
Phenotype of adult mouse muscle myoblasts reflects their fiber type of origin, 60:39-45

Fibronectin

Fehlner-Gardiner C.C., Uniyal S., von Ballestrem C.G., Chan B.M.C.
Differential utilization of VLA-4(α 4 β 1) and -5(α 5 β 1) integrins during the development of mouse bone marrow-derived mast cells, 60:317-325

Fin regeneration

Mari-Beffa M., Fernandez-Llebrez J.A.S.P., Becerra J.
Histochemically defined cell states during tail fin regeneration in teleost fishes, 60:139-149

Fish (see Fin)

FP receptor

Casimir D.A., Wilson-Miller C., Ntambi J.M.
Preadipocyte differentiation blocked by prostaglandin stimulation of prostanoid FP² receptor in murine 3T3-L1 cells, 60:203-210

4f-rnp gene

Hess, K.A., Simone A.A., Petschek J.P.
Spatial and temporal expression of 4f-rnp gene in *Drosophila melanogaster*, 61:103-111

Gap junctions

Matic M., Petrov I.N., Chen S., Wang C., Dimitrijevic S.D., Wolosin J.M.
Stem cells of the corneal epithelium lack connexins and metabolite transfer capacity, 61:251-260

Gastrulation

Vafa O., Goetzl L., Poccia D., Nishioka D.
Localization and characterization of blastocoelic extracellular matrix antigens in early sea urchin embryos and evidence for their proteolytic modification during gastrulation, 60:129-138

Gene expression

- Hermans-Borgmeyer I., Schinke B., Schaller H.C., Hoffmeister-Ullrich S.A.H.
Isolation of a marker for head-specific cell differentiation in hydra, 61:95-101
- Hess, K.A., Simone A.A., Petsehek J.P.
Spatial and temporal expression of 4f-rnp gene in *Drosophila melanogaster*, 61:103-111
- Itier J.-M., Douhet P., Desbois P., Joshi R.L., Dandoy-Dron F., Jami J., Bucchini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316
- Koide M., Akins R.E., Harayama H., Yasui K., Yokota M., Tuan R.S.
Atrial natriuretic peptide accelerates proliferation of chick embryonic cardiomyocytes in vitro, 61:1-11
- Liebert M., Hubbel A., Chung M., Wedemeyer G., Lomax M.I., Hegeman A., Yuan T.Y., Brozovich M., Wheelock M.J., Grossman H.B.
Expression of *mal* is associated with urothelial differentiation in vitro: identification by differential display reverse-transcriptase polymerase chain reaction, 61:177-185
- Rogers M.A., Langbein L., Praetzel S., Moll I., Krieg T., Winter H., Schweizer J.
Sequences and differential expression of three novel human type-II hair keratins, 61:187-194
- Yang C., Brar S.K., Desbarats L., Siu C.-H.
Synthesis of the Ca^{2+} -dependent cell adhesion molecule DdCAD-1 is regulated by multiple factors during Dictyostelium development, 61:275-284

Genomic imprinting

- McKarnay L.A., Overall M.L., Dziadek M.
Expression of H19 and Igf2 genes in uniparental mouse ES cells during in vitro and in vivo differentiation, 60:75-86

Germ cell cancer

- Moasser M.M., Khoo K.-S., Maerz W.J., Zelenetz A., Dmitrovsky E.
Derivation and characterization of retinoid-resistant human embryonal carcinoma cells, 60:251-257

Germ cells

- Moens A., Chastant S., Chesné P., Fléchon J.-E., Betteridge K.J., Renard J.-P.
Differential ability of male and female rabbit fetal germ cell nuclei to be reprogrammed by nuclear transfer, 60:339-345

Glucocorticoids

- Grigoriadis A.E., Heersche J.N.M., Aubin J.E.
Analysis of chondroprogenitor frequency and cartilage differentiation in a novel family of clonal chondrogenic rat cell lines, 60:299-307

Glucose-6-phosphatase

- Radaeva S., Bannasch P.
Changes in catalase and glucose-6-phosphatase distribution patterns within oval cell compartment as possible differentiation markers during viral hepatocarcinogenesis in woodchucks, 60:169-178

Golgi apparatus

- Chazaud B., Muriel M.-P., Aubery M., Decastel M.
Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Gonads

- Dorizzi M., Richard-Mercier N., Pieau C.
The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

- Fröjdman K., Pelliniemi I.J., Lendahl U., Virtanen I., Eriksson J.E.
The intermediate filament protein nestin occurs transiently in differentiating testis of rat and mouse, 61:243-249

- Levacher C., Gautier C., Saez J.-M., Habert R.
Immunohistochemical localization of transforming growth factor β 1 and β 2 in the fetal and neonatal rat ovary, 61:45-51

- Paranko J., Seitz J., Meinhardt A.
Developmental expression of heat shock protein 60 (HSP60) in the rat testis and ovary, 60:159-167

Growth inhibition

- Jing Y., Zhang J., Waxman S., Mira-y-Lopez R.
Upregulation of cytokeratins 8 and 18 in human breast cancer T47D cells is retinoid-specific and retinoic acid receptor-dependent, 60:109-117

H19

- McKarnay L.A., Overall M.L., Dziadek M.
Expression of H19 and Igf2 genes in uniparental mouse ES cells during in vitro and in vivo differentiation, 60:75-86

Hair formation

- Matzuzaki T., Inamatsu M., Yoshizato K.
The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297
- Rogers M.A., Langbein L., Praetzel S., Moll I., Krieg T., Winter H., Schweizer J.
Sequences and differential expression of three novel human type-II hair keratins, 61:187-194

Heat shock

- Koide M., Akins R.E., Harayama H., Yasui K., Yokota M., Tuan R.S.
Atrial natriuretic peptide accelerates proliferation of chick embryonic cardiomyocytes in vitro, 61:1-11

Heat shock protein (HSP60)

- Paranko J., Seitz J., Meinhardt A.
Developmental expression of heat shock protein 60 (HSP60) in the rat testis and ovary, 60:159-167

Heparan sulfate

- Toriyama K., Muramatsu H., Hoshino T., Torii S., Muramatsu T.
Evaluation of heparin-binding growth factors in rescuing morphogenesis of heparitinase-treated mouse embryonic lung explants, 61:161-167

Heparitinase

- Toriyama K., Muramatsu H., Hoshino T., Torii S., Muramatsu T.
Evaluation of heparin-binding growth factors in rescuing morphogenesis of heparitinase-treated mouse embryonic lung explants, 61:161-167

Hepatocytes

- Koike T., Shiojiri N.
Differentiation of the mouse hepatic primordium cultured in vitro, 61:35-43

Heterogeneity of tumors

- Anzai H., Kitada Y., Bucana C.D., Sanchez R., Omoto R., Fidler I.J.
Intratumoral heterogeneity and inverse correlation between expression of E-cadherin and collagenase type IV in human gastric carcinomas, 60:119-127

Hexamethylene bisacetamide

- Moasser M.M., Khoo K.-S., Maerz W.J., Zelenetz A., Dmitrovsky E.
Derivation and characterization of retinoid-resistant human embryonal carcinoma cells, 60:251-257

HIT insulinoma cells

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F., Jami J., Bucchini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316

HT-29 cells

Chazaud B., Muriel M.-P., Aubery M., Decastel M.
Organization of the endoplasmic reticulum-Golgi system is related to the state of enterocytic differentiation of human HT-29 cells, 60:179-191

Human insulin transcripts

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F., Jami J., Bucchini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316

Human muscle

Baroffio A., Hamann M., Bernheim L., Bochaton-Piallat M.-L., Gabbiani G., Bader C.R.
Identification of self-renewing myoblasts in the progeny of single human muscle satellite cells, 60:47-57

Hydra

Hermans-Borgmeyer I., Schinke B., Schaller H.C., Hoffmeister-Ullrich S.A.H.
Isolation of a marker for head-specific cell differentiation in hydra, 61:95-101

Hypostomal cell differentiation

Hermans-Borgmeyer I., Schinke B., Schaller H.C., Hoffmeister-Ullrich S.A.H.
Isolation of a marker for head-specific cell differentiation in hydra, 61:95-101

Igf2 gene

McKarney L.A., Overall M.L., Dziadek M.
Expression of H19 and Igf2 genes in uniparental mouse ES cells during in vitro and in vivo differentiation, 60:75-86

Immune system

Kadmon G., Altevogt P.
The cell adhesion molecule L1: species and cell-type-dependent multiple binding mechanisms, 61:143-150

Induction of differentiation

Lopashov G.V., Golubeva O.N., Zviadadze K.G.
Inductive capacity of living eye tissues from adult frogs, 61:237-242

Insulin gene

Itier J.-M., Douhet P., Desbois P., Joshi R.L., Dandoy-Dron F., Jami J., Bucchini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316

Insulin-like growth factors

Ernst C.W., McFarland D.C., White M.E.
Expression of insulin-like growth factor II (IGF-II), IGF binding protein-2 and myogenin during differentiation of myogenic satellite cells derived from the turkey, 61:25-33

Integrins

Breitkreutz D., Stark H.-J., Mirancea N., Tomakidi P., Steinbauer H., Fusenig N.E.
Integrin and basement membrane normalization in mouse grafts of human keratinocytes - Implications for epidermal homeostasis, 61:195-209

Fehlner-Gardiner C.C., Uniyal S., von Ballestrem C.G., Chan B.M.C.
Differential utilization of VLA-4($\alpha 4 \beta 1$) and -5($\alpha 5 \beta 1$) integrins during the development of mouse bone marrow-derived mast cells, 60:317-325

Kadmon G., Altevogt P.
The cell adhesion molecule L1: species and cell-type-dependent multiple binding mechanisms, 61:143-150

Intermediate filaments

Fröjdman K., Pelliniemi I.J., Lendahl U., Virtanen I., Eriksson J.E.
The intermediate filament protein nestin occurs transiently in differentiating testis of rat and mouse, 61:243-249

Plancha C.E.
Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Wroblewski J., Engström M., Edwall-Arvidsson C., Sjöberg G., Sejersen T., Lendahl U.
Distribution of nestin in the developing mouse limb bud in vivo and in micro-mass cultures of cells isolated from limb buds, 61:151-159

Intimal thickening

Slomp J., Gittenberger-de Groot A.C., Kotliansky V.E., Glukhova M.A., Bogers A.J.J.C., Poelman R.E.
Cytokeratin expression in human arteries pertinent to intimal thickening formation in the ductus arteriosus, 61:305-311

Invasion

Anzai H., Kitadai Y., Bucana C.D., Sanchez R., Omoto R., Fidler I.J.
Intratumoral heterogeneity and inverse correlation between expression of E-cadherin and collagenase type IV in human gastric carcinomas, 60:119-127

Islets of Langerhans

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F., Jami J., Bucchini D.
Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316
Yuan S., Rosenberg L., Paraskevas S., Agapitos D., Duguid W.P.
Transdifferentiation of human islets to pancreatic ductal cells in collagen matrix culture, 61:67-75

Junctional plaques

Paffenholz R., Franke W.W.
Identification and localization of a neuronally expressed member of the plakoglobin/armadillo multigene family, 61:293-304

Junctions

Schäfer S., Stumpp S., Franke W.W.
Immunological identification and characterization of the desmosomal cadherin Dsg2 in coupled and uncoupled epithelial cells and in human tissues, 60:99-108

Keratinocytes

Breitkreutz D., Stark H.-J., Mirancea N., Tomakidi P., Steinbauer H., Fusenig N.E.
Integrin and basement membrane normalization in mouse grafts of human keratinocytes - Implications for epidermal homeostasis, 61:195-209

Keratins

Bickenbach J.R., Longley M.A., Bundman D.S., Dominey A.M., Bowden P.E., Rothnagel A.M., Roop D.R.
A transgenic mouse model that recapitulates the clinical features of both neonatal and adult forms of the skin disease epidermolytic hyperkeratosis, 61:129-139

Jing Y., Zhang J., Waxman S., Mira-y-Lopez R.
Upregulation of cytokeratins 8 and 18 in human breast cancer T47D cells is retinoid-specific and retinoic acid receptor-dependent, 60:109-117

Moyer P.D., Kaufman A.H., Zhang Z., Kao C.W.-C., Spaulding A.G., Kao W.W.-Y.
Conjunctival epithelial cells can resurface denuded cornea, but do not transdifferentiate to express cornea-specific keratin 12 following removal of limbal epithelium in mouse, 60:31-38

Plancha C.E.
Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Rogers M.A., Langbein L., Praetzel S., Moll I., Krieg T., Winter H., Schweizer J.
Sequences and differential expression of three novel human type-II hair keratins, 61: 187-194

Laminin

Durham P.L., Snyder J.M.
Regulation of the β 2 subunit chain of laminin in developing rabbit fetal lung tissue, 60:229-243

LAMP-1

Cella N., Cornejo-Urbe R.R., Montes G.S., Hynes N.E.,
The lysosomal-associated membrane protein LAMP-1 is a novel differentiation marker for HC11 mouse mammary epithelial cells, 61:113-120

Larval antigen

Izutsu Y., Yoshizato K., Tochinal S.
Adult-type splenocytes of *Xenopus* induce apoptosis of histocompatible larval tail cells in vitro, 60:277-286

Lens

Okamoto M.
Simultaneous demonstration of lens regeneration from dorsal iris and tumour production from ventral iris in the same newt eye after carcinogen administration, 61:285-292

Wride M.A.
Cellular and molecular features of lens differentiation: a review of recent advances, 61:77-93

Letrozole

Dorizzi M., Richard-Mercier N., Pieau C.
The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

Lineages of chondrocytes

Grigoriadis A.E., Heersche J.N.M., Aubin J.E.
Analysis of chondroprogenitor frequency and cartilage differentiation in a novel family of clonal chondrogenic rat cell lines, 60:299-307

Liver primordium

Koike T., Shiojiri N.
Differentiation of the mouse hepatic primordium cultured in vitro, 61:35-43

Lung development

Durham P.L., Snyder J.M.
Regulation of the β 2 subunit chain of laminin in developing rabbit fetal lung tissue, 60:229-243

Toriyama K., Muramatsu H., Hoshino T., Torii S., Muramatsu T.
Evaluation of heparin-binding growth factors in rescuing morphogenesis of heparitinase-treated mouse embryonic lung explants, 61:161-167

Lymphocytes

Kadmon G., Altevogt P.
The cell adhesion molecule L1: species and cell-type-dependent multiple binding mechanisms, 61:143-150

Maki A., Mohammad R.M., Smith M., Al-Katib A.
Role of ubiquitin carboxyl terminal hydrolase in the differentiation of human acute lymphoblastic leukemia cell line, Reh, 60:59-66

Lysosomes

Cella N., Cornejo-Urbe R.R., Montes G.S., Hynes N.E.,
The lysosomal-associated membrane protein LAMP-1 is a novel differentiation marker for HC11 mouse mammary epithelial cells, 61:113-120

Mammary gland

Cella N., Cornejo-Urbe R.R., Montes G.S., Hynes N.E.,
The lysosomal-associated membrane protein LAMP-1 is a novel differentiation marker for HC11 mouse mammary epithelial cells, 61:113-120

Hayward S.W., Brody J.R., Cunha G.R.
An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219-227

Mast cells

Fehlner-Gardiner C.C., Uniyal S., von Ballestrem C.G., Chan B.M.C.
Differential utilization of VLA-4(α 4 β 1) and -5(α 5 β 1) integrins during the development of mouse bone marrow-derived mast cells, 60:317-325

Meiosis

Plancha C.E.
Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Merkel cells

Zhang C., Oakley B.
The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Metamorphosis

Izutsu Y., Yoshizato K., Tochinal S.
Adult-type splenocytes of *Xenopus* induce apoptosis of histocompatible larval tail cells in vitro, 60:277-286

Metastasis

Anzai H., Kitadai Y., Bucana C.D., Sanchez R., Omoto R., Fidler I.J.
Intratumoral heterogeneity and inverse correlation between expression of E-cadherin and collagenase type IV in human gastric carcinomas, 60:119-127

Morphogenesis

Toriyama K., Muramatsu H., Hoshino T., Torii S., Muramatsu T.
Evaluation of heparin-binding growth factors in rescuing morphogenesis of heparitinase-treated mouse embryonic lung explants, 61:161-167

Zilakos N.P., Zafiratos C.S., Parchment R.E.
Stage-dependent genetically-based deformities of the regenerating newt limb from 4-nitroquinoline-N-oxide mutagenesis: Potential embryonic regulation of cancer, 60:67-74

Muscle development

Eng H., Herrenknecht K., Semb H., Starzinski-Powitz A., Ringertz N., Gullberg D.
Effects of divalent cations on M-cadherins expression and distribution during primary rat myogenesis in vitro, 61:169-176

Ernst C.W., McFarland D.C., White M.E.

Expression of insulin-like growth factor II (IGF-II), IGF binding protein-2 and myogenin during differentiation of myogenic satellite cells derived from the turkey, 61:25-33

Franke W.W., Stehr S., Stump S., Kuhn C., Heid H., Rackwitz H.-R., Schnölzer M., Baumann R., Holzhausen H.-J., Moll R.

Specific immunohistochemical detection of cardiac/fetal α -actin in human cardiomyocytes and regenerating skeletal muscle cells, 60:245-250

Matschak T.W., Stickland N.C., Mason P.S., Crook A.R.

Oxygen availability and temperature affect embryonic muscle development in Atlantic salmon (*Salmo salar* L.), 61:229-235

Rosenblatt J.D., Parry D.J., Partridge T.A.

Phenotype of adult mouse muscle myoblasts reflects their fiber type of origin, 60:39-45

Wroblewski J., Engström M., Edwall-Arvidsson C., Sjöberg G., Sejersen T., Lendahl U.

Distribution of nestin in the developing mouse limb bud in vivo and in micro-mass cultures of cells isolated from limb buds, 61:151-159

Mutagenesis

Zilakos N.P., Zafiratos C.S., Parchment R.E.

Stage-dependent genetically-based deformities of the regenerating newt limb from 4-nitroquinoline-N-oxide mutagenesis: Potential embryonic regulation of cancer, 60:67-74

Myoblasts

Baroffio A., Hamann M., Bernheim L., Bochaton-Piallat M.-L., Gabbiani G., Bader C.R.

Identification of self-renewing myoblasts in the progeny of single human muscle satellite cells, 60:47-57

Rosenblatt J.D., Parry D.J., Partridge T.A.

Phenotype of adult mouse muscle myoblasts reflects their fiber type of origin, 60:39-45

Myogenin

Ernst C.W., McFarland D.C., White M.E.

Expression of insulin-like growth factor II (IGF-II), IGF binding protein-2 and myogenin during differentiation of myogenic satellite cells derived from the turkey, 61:25-33

Myosin heavy chain isoforms

Rosenblatt J.D., Parry D.J., Partridge T.A.

Phenotype of adult mouse muscle myoblasts reflects their fiber type of origin, 60:39-45

Myotubes

Rosenblatt J.D., Parry D.J., Partridge T.A.

Phenotype of adult mouse muscle myoblasts reflects their fiber type of origin, 60:39-45

Nervous system

Kadmon G., Altevogt P.

The cell adhesion molecule L1: species and cell-type-dependent multiple binding mechanisms, 61:143-150

Nestin

Fröjman K., Pelliniemi I.J., Lendahl U., Virtanen I., Eriksson J.E. The intermediate filament protein nestin occurs transiently in differentiating testis of rat and mouse, 61:243-249

Wroblewski J., Engström M., Edwall-Arvidsson C., Sjöberg G., Sejersen T., Lendahl U.

Distribution of nestin in the developing mouse limb bud in vivo and in micro-mass cultures of cells isolated from limb buds, 61:151-159

Neuronal proteins

Paffenholz R., Franke W.W.

Identification and localization of a neuronally expressed member of the plakoglobin/armadillo multigene family, 61:293-304

Nickel compounds

Okamoto M.

Simultaneous demonstration of lens regeneration from dorsal iris and tumour production from ventral iris in the same newt eye after carcinogen administration, 61:285-292

Nuclear bodies

Nason-Burchenal K.N., Maerz W., Albanell J., Allopenna J.,

Martin P., Moore M.A.S., Dmitrovsky E.

Common effects of different retinoic acid resistant promyelocytic leukemia cells are persistent telomerase activity and nuclear body disorganization, 61:321-331

Nuclear receptors

Wan Y.-J.Y., Wang L., Wu T.-C.J.

Murine endodermal F9E cells, derived from the teratocarcinoma line F9, contain high basal levels of retinoic acid receptors (RARs and RXRs) but are not sensitive to the actions of retinoic acid, 60:211-218

Nuclear transfer

Moens A., Chastant S., Chesné P., Fléchon J.-E., Betteridge K.J., Renard J.-P.

Differential ability of male and female rabbit fetal germ cell nuclei to be reprogrammed by nuclear transfer, 60:339-345

Ontogeny

Weber B., Rösel M., Arch R., Möller P., Zöller M.

Transient expression of CD44 variant isoforms in the ontogeny of the rat: Ectoderm-, endoderm- and mesoderm-derived cells express different exon combinations, 60:17-29

Oocyte maturation

Plancha C.E.

Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Oogenesis

Plancha C.E.

Cytokeratin dynamics during oocyte maturation in the hamster requires reaching of metaphase I, 60:87-98

Osteoblasts

Cohen-Tanugi A., Bolle M., Boy-Lefèvre M.-L., Anagnostou F., Forest N.

Phenotypic characterization of the 3A/1D-1M osteogenic cell line derived from in vivo transplantation of 3A/1D-1 chondroprogenitor murine teratocarcinoma cells, 60:327-337

Oval cells

Radaeva S., Bannasch P.

Changes in catalase and glucose-6-phosphatase distribution patterns within oval cell compartment as possible differentiation markers during viral hepatocarcinogenesis in woodchucks, 60:169-178

Ovarian masculinization

Dorizzi M., Richard-Mercier N., Pieau C.

The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

Ovary

Dorizzi M., Richard-Mercier N., Pieau C.

The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

Levacher C., Gautier C., Saez J.-M., Habert R.

Immunohistochemical localization of transforming growth factor $\beta 1$ and $\beta 2$ in the fetal and neonatal rat ovary, 61:45-51

Paranko J., Seitz J., Meinhardt A.

Developmental expression of heat shock protein 60 (HSP60) in the rat testis and ovary, 60:159-167

Pancreatic duct cells

Yuan S., Rosenberg L., Paraskevas S., Agapitos D., Duguid W.P.

Transdifferentiation of human islets to pancreatic ductal cells in collagen matrix culture, 61:67-75

3T3-L-1 preadipocyte

Casimir D.A., Wilson-Miller C., Ntambi J.M.

Preadipocyte differentiation blocked by prostaglandin stimulation of prostanoid FP_2 receptor in murine 3T3-L1 cells, 60:203-210

Ntambi J.M., Takova T.

Role of Ca^{2+} in the early stages of murine adipocyte differentiation as evidenced by calcium mobilizing agents, 60:151-158

Primordial germ cells

Moens A., Chastant S., Chesné P., Fléchon J.-E., Betteridge K.J., Renard J.-P.

Differential ability of male and female rabbit fetal germ cell nuclei to be reprogrammed by nuclear transfer, 60:339-345

Progenitors

Cohen-Tanugi A., Bolle M., Boy-Lefèvre M.-L., Anagnostou F., Forest N.

Phenotypic characterization of the 3A/1D-1M osteogenic cell line derived from in vivo transplantation of 3A/1D-1 chondroprogenitor murine teratocarcinoma cells, 60:327-337

Grigoriadis A.E., Heersche J.N.M., Aubin J.E.

Analysis of chondroprogenitor frequency and cartilage differentiation in a novel family of clonal chondrogenic rat cell lines, 60:299-307

Promoter analysis

Robinson V., Williams J.

A marker of terminal stalk cell terminal differentiation in *Dictyostelium*, 61:223-228

Promyelocytic leukemia

Nason-Burchenal K.N., Maerz W., Albanell J., Allopenna J.,

Martin P., Moore M.A.S., Dmitrovsky E.

Common effects of different retinoic acid resistant promyelocytic leukemia cells are persistent telomerase activity and nuclear body disorganization, 61:321-331

Proportion regulation

Schaap P., Tang Y., Othmer H.G.

A model for pattern formation in *Dictyostelium discoideum*, 60:1-16, 61:141

Prostaglandins

Casimir D.A., Wilson-Miller C., Ntambi J.M.

Preadipocyte differentiation blocked by prostaglandin stimulation of prostanoid FP_2 receptor in murine 3T3-L1 cells, 60:203-210

Prostate gland

Hayward S.W., Brody J.R., Cunha G.R.

An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219-227

Proteolysis

Vafa O., Goetzl L., Poccia D., Nishioka D.

Localization and characterization of blastocoelic extracellular matrix antigens in early sea urchin embryos and evidence for their proteolytic modification during gastrulation, 60:129-138

Quantitative modeling

Schaap P., Tang Y., Othmer H.G.

A model for pattern formation in *Dictyostelium discoideum*, 60:1-16, 61:141

Rabbit

Durham P.L., Snyder J.M.

Regulation of the $\beta 2$ subunit chain of laminin in developing rabbit fetal lung tissue, 60:229-243

Moens A., Chastant S., Chesné P., Fléchon J.-E., Betteridge K.J.,

Renard J.-P.

Differential ability of male and female rabbit fetal germ cell nuclei to be reprogrammed by nuclear transfer, 60:339-345

Rat

Eng H., Herrenknecht K., Semb H., Starzinski-Powitz A.,

Ringertz N., Gullberg D.

Effects of divalent cations on M-cadherins expression and distribution during primary rat myogenesis in vitro, 61:169-176

Fröjdman K., Pelliniemi I.J., Lendahl U., Virtanen I., Eriksson J.E. The intermediate filament protein nestin occurs transiently in differentiating testis of rat and mouse, 61:243-249

Grigoriadis A.E., Heersche J.N.M., Aubin J.E.

Analysis of chondroprogenitor frequency and cartilage differentiation in a novel family of clonal chondrogenic rat cell lines, 60:299-307

Hayward S.W., Brody J.R., Cunha G.R.

An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219-227

Levacher C., Gautier C., Saez J.-M., Habert R.

Immunohistochemical localization of transforming growth factor $\beta 1$ and $\beta 2$ in the fetal and neonatal rat ovary, 61:45-51

Matzuzaki T., Inamatsu M., Yoshizato K.

The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297

Weber B., Rösel M., Arch R., Möller P., Zöller M.

Transient expression of CD44 variant isoforms in the ontogeny of the rat: Ectoderm-, endoderm- and mesoderm-derived cells express different exon combinations, 60:17-29

Zhang C., Oakley B.

The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Regeneration

Franke W.W., Stehr S., Stump S., Kuhn C., Heid H.,

Rackwitz H.-R., Schnölzer M., Baumann R., Holzhausen H.-J.,

Moll R.

Specific immunohistochemical detection of cardiac/fetal α -actin in human cardiomyocytes and regenerating skeletal muscle cells, 60:245-250

Hermans-Borgmeyer I., Schinke B., Schaller H.C.,

Hoffmeister-Ullrich S.A.H.

Isolation of a marker for head-specific cell differentiation in hydra, 61:95-101

Marí-Beffa M., Fernandez-Llebrez J.A.S.P., Becerra J.

Histochemically defined cell states during tail fin regeneration in teleost fishes, 60:139-149

Matzuzaki T., Inamatsu M., Yoshizato K.

The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297

Okamoto M.

Simultaneous demonstration of lens regeneration from dorsal iris and tumour production from ventral iris in the same newt eye after carcinogen administration, 61:285-292

Reprogramming

Moens A., Chastant S., Chesné P., Fléchon J.-E., Betteridge K.J., Renard J.-P.

Differential ability of male and female rabbit fetal germ cell nuclei to be reprogrammed by nuclear transfer, 60:339-345

Restriction fragment length polymorphism

Moasser M.M., Khoo K.-S., Maerz W.J., Zelenetz A., Dmitrovsky E.
Derivation and characterization of retinoid-resistant human embryonal carcinoma cells, 60:251-257

Retinoic acid

Jing Y., Zhang J., Waxman S., Mira-y-Lopez R.
Upregulation of cytokeratins 8 and 18 in human breast cancer T47D cells is retinoid-specific and retinoic acid receptor-dependent, 60:109-117

Moasser M.M., Khoo K.-S., Maerz W.J., Zelenetz A., Dmitrovsky E.
Derivation and characterization of retinoid-resistant human embryonal carcinoma cells, 60:251-257

Naka K., Yokozaki H., Domen T., Hayashi K., Kuniyasu H., Yasui W., Lotan R., Tahara E.
Growth inhibition of cultured human gastric cancer cells by 9-*cis*-retinoic acid with induction of cdk inhibitor Waf1/Cip1/Sd1/p21 protein, 61:313-320

Nason-Burchenal K.N., Maerz W., Albanell J., Allopenna J., Martin P., Moore M.A.S., Dmitrovsky E.
Common effects of different retinoic acid resistant promyelocytic leukemia cells are persistent telomerase activity and nuclear body disorganization, 61:321-331

Wan Y.-J.Y., Wang L., Wu T.-C.J.
Murine endodermal F9E cells, derived from the teratocarcinoma line F9, contain high basal levels of retinoic acid receptors (RARs and RXRs) but are not sensitive to the actions of retinoic acid, 60:211-218

Retinoic acid receptors

Wan Y.-J.Y., Wang L., Wu T.-C.J.
Murine endodermal F9E cells, derived from the teratocarcinoma line F9, contain high basal levels of retinoic acid receptors (RARs and RXRs) but are not sensitive to the actions of retinoic acid, 60:211-218

Moasser M.M., Khoo K.-S., Maerz W.J., Zelenetz A., Dmitrovsky E.
Derivation and characterization of retinoid-resistant human embryonal carcinoma cells, 60:251-257

Retinoid resistance

Naka K., Yokozaki H., Domen T., Hayashi K., Kuniyasu H., Yasui W., Lotan R., Tahara E.
Growth inhibition of cultured human gastric cancer cells by 9-*cis*-retinoic acid with induction of cdk inhibitor Waf1/Cip1/Sd1/p21 protein, 61:313-320

Nason-Burchenal K.N., Maerz W., Albanell J., Allopenna J., Martin P., Moore M.A.S., Dmitrovsky E.
Common effects of different retinoic acid resistant promyelocytic leukemia cells are persistent telomerase activity and nuclear body disorganization, 61:321-331

Reverse-transcriptase polymerase chain reaction

Liebert M., Hubbel A., Chung M., Wedemeyer G., Lomax M.I., Hegeman A., Yuan T.Y., Brozovich M., Wheelock M.J., Grossman H.B.
Expression of *mal* is associated with urothelial differentiation in vitro: identification by differential display reverse-transcriptase polymerase chain reaction, 61:177-185

RNA binding proteins

Hess, K.A., Simone A.A., Petschek J.P.
Spatial and temporal expression of 4f-rnp gene in *Drosophila melanogaster*, 61:103-111

RNA editing

Hess, K.A., Simone A.A., Petschek J.P.
Spatial and temporal expression of 4f-rnp gene in *Drosophila melanogaster*, 61:103-111

Salivary gland

Hayward S.W., Brody J.R., Cunha G.R.
An edgewise look at basal epithelial cells: Three-dimensional views of the rat prostate, mammary gland and salivary gland, 60:219-227

Salmon

Matschak T.W., Stickland N.C., Mason P.S., Crook A.R.
Oxygen availability and temperature affect embryonic muscle development in Atlantic salmon (*Salmo salar* L.), 61:229-235

Satellite cells

Baroffio A., Hamann M., Bernheim L., Bochaton-Piallat M.-L., Gabbiani G., Bader C.R.
Identification of self-renewing myoblasts in the progeny of single human muscle satellite cells, 60:47-57

Ernst C.W., McFarland D.C., White M.E.
Expression of insuline-like growth factor II (IGF-II), IGF binding protein-2 and myogenin during differentiation of myogenic satellite cells derived from the turkey, 61:25-33

Franke W.W., Stehr S., Stumpp S., Kuhn C., Heid H., Rackwitz H.-R., Schnölzer M., Baumann R., Holzhausen H.-J., Moll R.
Specific immunohistochemical detection of cardiac/fetal α -actin in human cardiomyocytes and regenerating skeletal muscle cells, 60:245-250

Rosenblatt J.D., Parry D.J., Partridge T.A.
Phenotype of adult mouse muscle myoblasts reflects their fiber type of origin, 60:39-45

Sea urchin

Vafa O., Goetzl L., Poccia D., Nishioka D.
Localization and characterization of blastocoelic extracellular matrix antigens in early sea urchin embryos and evidence for their proteolytic modification during gastrulation, 60:129-138

Signal transduction

Kadmon G., Altevogt P.
The cell adhesion molecule L1: species and cell-type-dependent multiple binding mechanisms, 61:143-150

Signaling network

Schaap P., Tang Y., Othmer H.G.
A model for pattern formation in *Dictyostelium discoideum*, 60:1-16, 61:141

Smooth muscle cells

Slomp J., Gittenberger-de Groot A.C., Kotliansky V.E., Glukhova M.A., Bogers A.J.J.C., Poelman R.E.
Cytokeratin expression in human arteries pertinent to intimal thickening formation in the ductus arteriosus, 61:305-311

Splenic leukocytes

Izutsu Y., Yoshizato K., Tochinali S.
Adult-type splenocytes of *Xenopus* induce apoptosis of histocompatible larval tail cells in vitro, 60:277-286

staB

Robinson V., Williams J.
A marker of terminal stalk cell terminal differentiation in *Dictyostelium*, 61:223-228

Stalk cells

Robinson V., Williams J.
A marker of terminal stalk cell terminal differentiation in *Dictyostelium*, 61:223-228

Stearoyl-CoA desaturase

Ntambi J.M., Takova T.
Role of Ca^{2+} in the early stages of murine adipocyte differentiation as evidenced by calcium mobilizing agents, 60:151-158

Stem cells

Baroffio A., Hamann M., Bernheim L., Bochaton-Piallat M.-L., Gabbiani G., Bader C.R.
Identification of self-renewing myoblasts in the progeny of single human muscle satellite cells, 60:47-57

Henderson K., Kirkland S.C.
Multilineage differentiation of cloned HRA-19 cells in serum-free medium: A model of human colorectal epithelial differentiation, 60:259-268

Lin Y.I.
Day 11 mouse fetal thymus: phenotype and search for the point of commitment, 61:53-65

Matic M., Petrov I.N., Chen S., Wang C., Dimitrijevic S.D., Wolosin J.M.
Stem cells of the corneal epithelium lack connexins and metabolite transfer capacity, 61:251-260

Matuzaki T., Inamatsu M., Yoshizato K.
The upper dermal sheath has a potential to regenerate the hair in the rat follicular epidermis, 60:287-297

McKarney L.A., Overall M.L., Dziadek M.
Expression of H19 and Igf2 genes in uniparental mouse ES cells during in vitro and in vivo differentiation, 60:75-86

Taste buds

Zhang C., Oakley B.
The distribution and origin of keratin 20-containing taste buds in rat and human, 61:121-127

Telomerase

Nason-Burchenal K.N., Maerz W., Albanell J., Allopenna J., Martin P., Moore M.A.S., Dmitrovsky E.
Common effects of different retinoic acid resistant promyelocytic leukemia cells are persistent telomerase activity and nuclear body disorganization, 61:321-331

Temperature-dependent development

Matschak T.W., Stickland N.C., Mason P.S., Crook A.R.
Oxygen availability and temperature affect embryonic muscle development in Atlantic salmon (*Salmo salar* L.), 61:229-235

Temperature-dependent sex determination

Dorizzi M., Richard-Mercier N., Pieau C.
The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

Teratocarcinoma cells

Cohen-Tanugi A., Bolle M., Boy-Lefèvre M.-L., Anagnostou F., Forest N.
Phenotypic characterization of the 3A/1D-1M osteogenic cell line derived from in vivo transplantation of 3A/1D-1 chondroprogenitor murine teratocarcinoma cells, 60:327-337

Wan Y.-J.Y., Wang L., Wu T.-C.J.
Murine endodermal F9E cells, derived from the teratocarcinoma line F9, contain high basal levels of retinoic acid receptors (RARs and RXRs) but are not sensitive to the actions of retinoic acid, 60:211-218

Teratomas

McKarney L.A., Overall M.L., Dziadek M.
Expression of H19 and Igf2 genes in uniparental mouse ES cells during in vitro and in vivo differentiation, 60:75-86

Testis

Fröjdman K., Pelliniemi I.J., Lendahl U., Virtanen I., Eriksson J.E.
The intermediate filament protein nestin occurs transiently in differentiating testis of rat and mouse, 61:243-249

Paranko J., Seitz J., Meinhardt A.
Developmental expression of heat shock protein 60 (HSP60) in the rat testis and ovary, 60:159-167

TGF β

Levacher C., Gautier C., Saez J.-M., Habert R.
Immunohistochemical localization of transforming growth factor β 1 and β 2 in the fetal and neonatal rat ovary, 61:45-51

Thermosensitive period

Dorizzi M., Richard-Mercier N., Pieau C.
The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

Thymus

Lin Y.I.
Day 11 mouse fetal thymus: phenotype and search for the point of commitment, 61:53-65

Tight junctions

Liebert M., Hubbel A., Chung M., Wedemeyer G., Lomax M.I., Hegeman A., Yuan T.Y., Brozovich M., Wheelock M.J., Grossman H.B.
Expression of *mal* is associated with urothelial differentiation in vitro: identification by differential display reverse-transcriptase polymerase chain reaction, 61:177-185

Ti-plasmid

Meyer A.D., Aebi R., Meins Jr. F.
Tobacco plants carrying a *tms* locus of Ti-plasmid origin and the *HI-1* allele are tumor prone, 61:213-221

Tissue interaction

Koike T., Shiojiri N.
Differentiation of the mouse hepatic primordium cultured in vitro, 61:35-43

Tobacco plant

Meyer A.D., Aebi R., Meins Jr. F.
Tobacco plants carrying a *tms* locus of Ti-plasmid origin and the *HI-1* allele are tumor prone, 61:213-221

Transdifferentiation

Moyer P.D., Kaufman A.H., Zhang Z., Kao C.W.-C., Spaulding A.G., Kao W.W.-Y.
Conjunctival epithelial cells can resurface denuded cornea, but do not transdifferentiate to express cornea-specific keratin 12 following removal of limbal epithelium in mouse, 60:31-38

Yuan S., Rosenberg L., Paraskevas S., Agapitos D., Duguid W.P.
Transdifferentiation of human islets to pancreatic ductal cells in collagen matrix culture, 61:67-75

Transformation

Meyer A.D., Aebi R., Meins Jr. F.
Tobacco plants carrying a *tms* locus of Ti-plasmid origin and the *HI-1* allele are tumor prone, 61:213-221

Transforming growth factors

Levacher C., Gautier C., Saez J.-M., Habert R.
Immunohistochemical localization of transforming growth factor β 1 and β 2 in the fetal and neonatal rat ovary, 61:45-51

Transgenic insulin

Itier J.-M., Douhet P., Desbois P., Joshi R.-L., Dandoy-Dron F., Jami J., Bucchini D.

Human insulin gene expression in transgenic mice: mutational analysis of the regulatory region, 60:309-316

Trichocytes

Rogers M.A., Langbein L., Praetzel S., Moll I., Krieg T., Winter H., Schweizer J.

Sequences and differential expression of three novel human type-II hair keratins, 61: 187-194

Turkey

Ernst C.W., McFarland D.C., White M.E.

Expression of insuline-like growth factor II (IGF-II), IGF binding protein-2 and myogenin during differentiation of myogenic satellite cells derived from the turkey, 61:25-33

Turtle

Dorizzi M., Richard-Mercier N., Pieau C.

The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*, 60:193-201

Urothelial cells

Liebert M., Hubbel A., Chung M., Wedemeyer G., Lomax M.I., Hegeman A., Yuan T.Y., Brozovich M., Wheelock M.J., Grossman H.B.

Expression of *mal* is associated with urothelial differentiation in vitro: identification by differential display reverse-transcriptase polymerase chain reaction, 61:177-185

Vacuolar apical compartment

Lincz L.F., Buret A., Burns G.F.

Formation of spheroid structures in a human colon carcinoma cell line involves a complex series of intercellular rearrangements, 61:216-274

Viral hepatocarcinogenesis

Radaeva S., Bannasch P.

Changes in catalase and glucose-6-phosphatase distribution patterns within oval cell compartment as possible differentiation markers during viral hepatocarcinogenesis in woodchucks, 60:169-178

Wound-healing

Moyer P.D., Kaufman A.H., Zhang Z., Kao C.W.-C., Spaulding A.G., Kao W.W.-Y.

Conjunctival epithelial cells can resurface denuded cornea, but do not transdifferentiate to express cornea-specific keratin 12 following removal of limbal epithelium in mouse, 60:31-38

Xenopus laevis

Izutsu Y., Yoshizato K., Tochinal S.

Adult-type splenocytes of *Xenopus* induce apoptosis of histocompatible larval tail cells in vitro, 60:277-286

Lopashov G.V., Golubeva O.N., Zviadadze K.G.

Inductive capacity of living eye tissues from adult frogs, 61:237-242

Zebrafish

Marí-Beffa M., Fernandez-Llebrez J.A.S.P., Becerra J.

Histochemically defined cell states during tail fin regeneration in teleost fishes, 60:139-149